

ANAPHYLAXIS EMERGENCY ACTION PLAN

| | ZENT NATIE: | | AGE: | |
|--|--|----------------------------------|--|--|
| | RGY TO: | | | |
| ASTH | IMA: Tes (high risk for sever | e reaction) 🗆 NO | | |
| Oth | er health problems besides anaph | ylaxis: | | |
| Curi | rent medications, if any: | | | |
| Wea | ar medical identication jewelry th | at identifies the anaphylaxis po | otential and the food allergen triggers. | |
| YMP | TOMS OF ANAPHYLAXIS | INCLUDE: | | |
| • M | OUTH—itching, swelling of lips and/ | or tongue • Gl | JT—vomiting, diarrhea, cramps | |
| • THROAT*—itching, tightness/closure, hoarseness | | hoarseness • LU | LUNG*—shortness of breath, cough, wheeze | |
| • SK | (IN—itching, hives, redness, swelling | • HE | ART*—weak pulse, dizziness, passing out | |
| | nly a few symptoms may be present. Some symptoms can be life-threaten | | e quickly. | |
| ИНА Т | г то до: | | | |
| ١. | INJECT EPINEPHRINE IN TH | GH USING (check one): | | |
| | ☐ Adrenaclick (0.15 mg) | · | piPen Jr (0.15 mg) | |
| | □ Adrenaclick (0.30 mg) | | piPen (0.30 mg) | |
| | | a prescription for th eright med | epinephrine; medications shown in alpha order; ication for this patient, that it is current/not expired | |
| | Other medication/dose/route: | | | |
| IMI | PORTANT: Asthma inhale | rs and/or antihistamine | s can't be depended on in anaphylaxis | |
| | CALL 9-1-1 or RESCUE SQUA | AD (before calling contacts)! | ! | |
| 2. | EMEDICINICY CONTACTS | | | |
| | EMERGENCY CONTACTS | | | |
| 3. | #I home | work | cell | |
| 3. | | | | |

STUDENT'S SIGNATURE/DATE